

BEELOUSE, BRAULA COECA NITZSCH (DIPTERA: BRAULIDAE)¹

H. V. Weems, Jr.²

INTRODUCTION: The beelouse, Braula coeca Nitzsch 1818 (fig. 1), is a tiny commensal wingless fly found in colonies of the honey bee, Apis mellifera Linnaeus, where it lives on the bodies of the bees and literally steals its food out of the mouth of its host. As many as 187 beelice have been reported on a single bee queen. The beelouse was first mentioned in European literature in 1740 by Reaumur, who discussed its relation to the honey bee. Skaife (1921) and Argo (1927) described fairly completely the life cycle of the insect, while Arnhart (1923) found the mines and dried out larval skins under honey cappings. The beelouse is found on worker, drone, and queen honey bees (Smith 1978). It probably was first introduced into the United States from Europe along with the first shipments of honey bees. Some writers have stated that the beelouse causes little or no harm to bee colonies, but most investigators think that Braula is harmful. Most beekeepers practice mechanical control unknowingly by extracting honey because the Braula larvae are eliminated while removing cappings before extraction. Chemical control measures have been developed in Europe and Asia, where the beelouse is much more common than in North America, but no pesticide has been registered for use against the beelouse in the United States. Because of the frequent movement of large numbers of bee colonies throughout much of the United States, there probably is little that can be done to prevent the spread of the beelouse, and its control may be accomplished through mechanical practices of the beekeepers. However, Florida beekeepers should be alert to the occurrence of the beelouse.

DISTRIBUTION: This fly has been reported in Africa from Congo, Egypt, and Morocco, in Asia from India and the Soviet Union, in Australia from Tasmania, in most of Europe, in South America from Argentina, Brasil, Trinidad and Tobago, and Venezuela, and in North America from the United States: Alabama, Delaware, Illinois, Maryland, Minnesota, New York, North Carolina, Ohio, Pennsylvania, Tennessee, Virginia, West Virginia, and Wisconsin. In Florida it is known only from a single specimen collected on a queen honey bee by a DPI Apiary Inspector, James Hall, in Lake County near Umatilla on 17 March 1983.

IDENTIFICATION: Eggs are white, oval-shaped, with 2 lateral flanges which are flat and extend parallel to each other and to the long axis of the egg. Imms (1942) reported that the average length, without flanges, ranged from 0.78-0.81 mm and the width from 0.28-0.33 mm. Including the flanges, a typical egg measured 0.84 mm by 0.42 mm. Eggs may be deposited in various places in a hive--in empty cells, on brood cell cappings, or on wax dirt on the floor of a colony, but only eggs that are laid on honey cappings hatch. Incubation period for eggs varies from 2 days during summer to 7.4 days during winter. Larvae emerge from the attached end of their egg where they begin constructing a tunnel under cappings and sometimes on the walls and bottoms of cells. These tunnels give cappings of infested comb the appearance of being intersected with fine fractures, similar to the mines of a leaf miner. Larvae feed upon honey and pollen grains within the wax of their tunnels. Beelouse larvae pass through 3 instars, requiring from 7.1-10.8 days to complete larval development, depending upon the season of the year. The prepupa ranges from 1-2.7 days and appears cream-white when viewed through the transparent larval skin. The pupa is enclosed within the unmodified cuticle of the last larval instar and is white or yellowish in color, 1.4-1.7 mm long by 0.5-0.75 mm wide. The beelouse has eye rudi-

¹Contribution No. 566, Bureau of Entomology.

²Taxonomic Entomologist, P. O. Box 1269, Gainesville, FL 32602.

ments just above the antennae seen as pale spots on the cuticle surface surrounded by more darkly pigmented chitinous rings. There is no trace of wings or halteres. The tarsi are 5 segmented; each terminal joint contains a comb-like structure, divided in the middle, with a variable number of teeth (fig. 2). The combs allow Braula to cling tightly to the host.

SURVEY AND DETECTION: Careful inspection of individual bees within the hive is the only effective method of collecting specimens of the tiny beelouse. Specimens may be preserved in 75% isopropyl alcohol.

REFERENCES:

- Argo, V. N. 1927. Braula coeca in den Vereinigten Staaten. Archiv für Bienenkunde 8(1):48-52.
- Arnhart, L. 1923. Zur Entwicklungsgeschichte der Braula coeca Nitzsch. Zool. Anz. 56(9-10):193-97, 1 fig.
- Imms, A. D. 1942. On Braula coeca Nitzsch and its affinities. Parasit. 34:88-100.
- Nitzsch, C. L. 1818. Die Familien und Gattungen der Thierinsekten (Insecta epizoi- ca); als prodromus einer naturgeschichte der selben. Magazin Ent. 3:261-316.
- Reaumur, R.A.F. de. 1740. Memoirs pour servir a l'histoire des insectes, 5. Paris.
- Skaife, S. H. 1921. The life-history of the bee louse. S. African Bee. J. 1:38-9.
- Smith, I. B., Jr. 1978. The bee louse, Braula coeca Nitzsch, its distribution and biology on honey bees. Unpub. MS thesis, Univ. Maryland. 111 p., 6 fig., 12 tab., appendix.



FIGURES: 1. Frontal view of the beelouse, Braula coeca L.; 2. Dorsal view of a tarsus showing the comb-shaped row of spines, or claw, important in clinging to the branched hairs of the bee. (Scanning electron microscope photographs, compliments of Dr. W. J. Humphreys, University of Georgia).

